# The conformable condition for Nanodiscs

# Mariana M. F. da Cruz<sup>1</sup>, Celina M. H. de Figueiredo<sup>1</sup>, Diana Sasaki<sup>2</sup>, Marcus T. Costa<sup>2</sup>

<sup>1</sup>Programa de Engenharia de Sistemas e Computação (PESC) Rio de Janeiro Federal University (UFRJ) – Cidade Universitária, Rio de Janeiro – Brazil

<sup>2</sup>Instituto de Matemática e Estatística (IME) Rio de Janeiro State University (UERJ) – Maracanã, Rio de Janeiro – Brazil

{mmartins,celina}@cos.ufrj.br, {diana.sasaki,marcus.tovar}@ime.uerj.br

**Abstract.** We investigate the total coloring of fullerene nanodiscs, a subclass of cubic planar graphs with girth 5 arising in Chemistry, motivated by a conjecture about the nonexistence of a Type 2 cubic graph of girth at least 5. We give a combinatorial description and then a conformable coloring for an infinite family of fullerene nanodiscs.

# 1. The large girth Type 1 conjecture

A k-total coloring of a graph G is a color assignment from set  $E \cup V$ , where E denotes the set of edges and V denotes the set of vertices of the graph, such that distinct colors are assigned to: every pair of vertices that are adjacent; all edges that are adjacent; and each vertex and its incident edges. The total chromatic number  $\chi''(G)$  is the smallest natural k for which G admits a k-total coloring. Behzad and Vizing [Behzad 1965, Vizing 1964] independently conjectured the Total Coloring Conjecture (TCC) that for any simple graph G, we have  $\chi''(G) \leq \Delta(G) + 2$ . If  $\chi''(G) = \Delta(G) + 1$ , then the graph is Type 1; if  $\chi''(G) = \Delta(G) + 2$ , then the graph is Type 2. The TCC has been verified for some particular classes of graphs, including cubic graphs [Kostochka 1996].




Figure 1. (a) A Type 2 girth 3 cubic graph; (b) a Type 2 girth 4 cubic graph.

Every known Type 2 cubic graph has a girth 3 or 4 (See Figure 1). So, it is natural to think that there are no Type 2 cubic graphs with girth at least 5. Thus the following conjecture was proposed [Brinkmann et al. 2015]:

#### Conjecture 1. There is no Type 2 cubic graph with girth at least 5.

We will then define a special vertex coloring which is a necessary condition for a graph to be Type 1.

**Lemma 1** ([Chetwynd and Hilton 1988]). Let G be a regular graph. G is conformable if and only if it has a vertex coloring with  $\Delta + 1$  colors and each color class has the same parity of |V(G)|.

By Lemma 1, a necessary step towards proving that a cubic graph is Type 1 is to define a 4-vertex coloring where the cardinality of each vertex color class is even. Deciding whether an arbitrary graph is Type 1 is NP-complete [Sánchez-Arroyo 1989], even restricted to cubic graphs. Therefore, extending a conformable coloring to a  $(\Delta + 1)$ -total coloring is also NP-complete.

Lemma 2 ([Chetwynd and Hilton 1988]). If G is a Type 1 graph, then G is conformable.

## 2. The nanodiscs are graphs of girth 5

The fullerene nanodiscs, or nanodiscs  $D_r$  of radius  $r \ge 2$ , are structures composed of two identical flat covers, made with only hexagonal faces, connected by a strip along their borders constructed with additional 12 pentagonal faces. Figure 2 shows nanodiscs, where we highlighted with blue color in the connecting strip the 12 pentagonal faces.

The sequence  $\{1, 6, 12, 18, ..., 6(r-1), 6r, 6(r-1), ..., 18, 12, 6, 1\}$  provides the amount of faces on each layer of the nanodisc graph  $D_r$ . The 12 pentagonal faces are distributed in the central layer among its 6r faces with the other (6r - 12) hexagonal faces [Nicodemos 2017]. The auxiliary cycle sequence  $\{C_6, C_{18}, ..., C_{12r-6}, C_{12r-6}, ..., C_{18}, C_6\}$  provides the sizes of the auxiliary cycles that define the layers. A nanodisc contains  $12r^2$  vertices,  $18r^2$  edges and has girth 5.

## 3. All even radius nanodiscs are conformable

A strategy to color the vertices of  $D_r$  is to take advantage that the auxiliary cycles have even length and color alternately with colors 1 and 2 the cycle  $C_6$  defining the inner layer, with colors 3 and 4 the next cycle  $C_{18}$ , and so on. The strategy does not rely on the unicity of  $D_r$ , and defines for even radius a 4-vertex coloring that is conformable. See Figure 2.




Figure 2. Conformable colorings for  $D_2$  and  $D_4$ , respectively.

**Theorem 1** ([da Cruz et al. 2021]). *Every nanodisc with even radius admits a conformable coloring.* 

*Proof.* In a  $D_r$  with even r, consider the 4-vertex coloring that gives colors 1 and 2 to the outer cycle  $C_6$ , colors 3 and 4 to the next cycle  $C_{18}$ , until we reach the central layer, where colors 3 and 4 are given to cycle  $C_{12r-6}$  and colors 1 and 2 are given to the next cycle  $C_{12r-6}$ , continue in this fashion until colors 3 and 4 are given to the inner cycle  $C_6$ . There are 2r cycles, and each color appears in r cycles. Each color class has the same even number  $3r^2$  of vertices.




Figure 3. (a) A 4-vertex coloring that does not give a conformable coloring of  $D_3$ ; (b) an optimal 3-vertex coloring that gives a conformable coloring of  $D_3$ .

Note that for r odd this 4-vertex coloring is not conformable, since  $3r^2$  generates an odd number of vertices for each color class (See Figure 3(a)). Thus, seeking to prove that all  $D_r$  nanodiscs are conformable, we further studied the  $D_3$  nanodisc and obtained an optimal 3-vertex coloring that gives a conformable 4-vertex coloring, since each of the three color classes has an even number of vertices, and the fourth color class has 0 elements (See Figure 3(b)).

**Theorem 2** ([da Cruz et al. 2021]). *The nanodisc*  $D_3$  *is conformable.* 




Figure 4. 3-vertex colorings of the two non-isomorphic instances of  $D_5$ .

The optimal 3-vertex coloring strategy for  $D_3$  consists of coloring the auxiliary cycles, except for the two cycles of the center layer, using two colors alternately, avoiding color conflict in the vertices that are extremes of radial edges between consecutive auxiliary cycles. In the cycles  $C_{12r-6}$  defining the central layer, we introduce a third color by choosing six vertices in each cycle, thus ensuring the parity of this color class and the other color classes, where the fourth color class has 0 vertices. Recently, we have found two non-isomorphic instances for the fullerene nanodisc  $D_5$ . We verified the coloring strategy described above in both, which culminated in the following result (See Figure 4).

**Theorem 3.** The two non-isomorphic instances of nanodisc  $D_5$  are conformable.

#### 4. Current work

Our current goal is to generalize the 3-vertex coloring strategy that gives a conformable coloring of  $D_3$  presented in Figure 3(b) and of  $D_5$  presented in Figure 4 to an arbitrary nanodisc of odd radius. We were not able to extend the conformable vertex colorings defined in Section 3 for a 4-total coloring of those fullerene nanodiscs. We need to further investigate the structure of the class, looking for a suitable conformable vertex coloring that extends to a 4-total coloring of  $D_r$ , in order to prove that every fullerene nanodisc is Type 1, as has already been proven for  $D_2$  [da Cruz et al. 2021]. See Figure 5.




Figure 5. A 4-total coloring of  $D_2$ .

#### References

Behzad, M. (1965). Graphs and their chromatic numbers. Michigan State University.

- Brinkmann, G., Preissmann, M., and Sasaki, D. (2015). Snarks with total chromatic number 5. *Discrete Mathematics and Theoretical Computer Science*, 17:369–382.
- Chetwynd, A. and Hilton, A. (1988). Some refinements of the total chromatic number conjecture. *Congressus Numerantium*, 66:195–216.
- da Cruz, M., de Figueiredo, C., Sasaki, D., and Costa, M. (2021). Hunting a Type 2 fullerene nanodisc. *Matemática Contemporânea*, 48:126–136.
- Kostochka, A. V. (1996). The total chromatic number of any multigraph with maximum degree five is at most seven. *Discrete Mathematics*, 162:199–214.
- Nicodemos, D. (2017). Diâmetro de Grafos Fulerenes e Transversalidade de Ciclos Ímpares de Fuleróides-(3, 4, 5, 6). PhD thesis, Universidade Federal do Rio de Janeiro.
- Sánchez-Arroyo, A. (1989). Determining the total colouring number is NP-hard. *Discrete Mathematics*, 78(3):315–319.
- Vizing, V. (1964). On an estimate of the chromatic class of a p-graph. *Discret Analiz*, 3:25–30.