# On the maximum number of edges in a graph with prescribed walk-nonrepetitive chromatic number \*

Fábio Botler<sup>1</sup>, Wanderson Lomenha<sup>1</sup>, João Pedro de Souza<sup>1,2</sup>

<sup>1</sup>Programa de Engenharia de Sistemas e Computação Instituto Alberto Luiz Coimbra de Pós-Graduação e Pesquisa em Engenharia Universidade Federal do Rio de Janeiro, Brasil

> <sup>2</sup>Departamento de Matemática Colégio Pedro II

{fbotler, wlomenha, jpsouza}@cos.ufrj.br

Abstract. Fix a coloring  $c: V(G) \to \mathbb{N}$  of the vertices of a graph G and let  $W = v_1 \cdots v_{2r}$  be a walk in G. We say that W is repetitive (with respect to c) if  $c(v_i) = c(v_{i+r})$  for every  $i \in \{1, \ldots, r\}$ ; and that W is boring if  $v_i = v_{i+r}$ , for every  $i \in \{1, \ldots, r\}$ . Finally, we say that c is a walk-nonrepetitive coloring of G if every repetitive walk is boring, and we denote by  $\sigma(G)$  the walk-nonrepetitive chromatic number, i.e., the minimum number of colors in a walk-nonrepetitive coloring of G. In this paper we explore the maximum number of edges in a graph G with n vertices for which  $\sigma(G) = k$ , for  $k \ge 4$ . In [Barát and Wood 2008] it was shown that  $e(G) \le \frac{1}{2}(k-1)n$ . We show that  $e(G) = \frac{1}{2}(k-1)n$  if and only if G is a union of disjoint copies of  $K_k$ . We also show that this upper bound can be improved for connected graphs for the case k = 4: if G is a connected graph for which  $\sigma(G) \ge 5$ , then  $e(G) \le \frac{4}{3}|V(G)|$ .

**Resumo.** Fixe uma coloração  $c: V(G) \to \mathbb{N}$  dos vértices de G e seja  $W = v_1 \cdots v_{2r}$  um passeio em G. Dizemos que W é repetitivo (com respeito a c) se  $c(v_i) = c(v_{i+r})$  para todo  $i \in \{1, \ldots, r\}$ ; e que W é entendiante se  $v_i = v_{i+r}$  para todo todo  $i \in \{1, \ldots, r\}$ . Finalmente, dizemos que c é uma coloração passeio não-repetitiva de G se todo passeio repetitivo é entendiante, e denotamos por  $\sigma(G)$  o número cromático passeio não-repetitivo, *i.e.*, o menor número de cores em uma coloração passeio não-repetitiva de G. Neste artigo, exploramos o número máximo de arestas em um grafo G com n vértices para o qual  $\sigma(G) = k$ , para  $k \ge 4$ . Em [Barát and Wood 2008] foi provado que  $e(G) \le \frac{1}{2}(k-1)n$ . Nós mostramos que  $e(G) = \frac{1}{2}(k-1)n$  se e somente se G é uma união de cópias disjuntas de  $K_k$ . Também provamos que o limitante superior pode ser melhorado para grafos conexos quando k = 4: se G é um grafo conexo com  $\sigma(G) = 4 e |V(G)| \ge 5$ , então  $e(G) \le \frac{4}{3}|V(G)|$ .

## 1. Introduction

All graphs in this paper are finite, undirected and simple. Given a graph G and a vertex  $v \in V(G)$ , we denote by d(v) (resp. N(v)) the *degree* (resp. the set of neighbors) of v.

<sup>\*</sup>This research has been partially supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil – CAPES – Finance Code 001. F. Botler is partially supported by CNPq (423395/2018-1), FAPERJ (211.305/2019) and UFRJ (23733). W. Lomenha is partially supported by CNPq (140654/2021-6). FAPERJ is the Rio de Janeiro Research Foundation. CNPq is the National Council for Scientific and Technological Development of Brazil.

A walk in G is a sequence  $v_1 \cdots v_r$  of vertices of G for which  $v_i v_{i+1} \in E(G)$  for every  $i \in \{1, \ldots, r-1\}$ . A walk  $W = v_1 \cdots v_{2r}$  is boring if  $v_i = v_{i+r}$  for each  $i \in \{1, \ldots, r\}$ . Given a coloring  $c \colon V(G) \to \mathbb{N}$ , we say that W is repetitive if  $c(v_i) = c(v_{i+r})$  for each  $i \in \{1, \ldots, r\}$ . Finally, we say that a coloring  $c \colon V(G) \to \{1, \ldots, k\}$  is walk-nonrepetitive if every repetitive walk is boring, and denote by  $\sigma(G)$  the minimum k for which there is a walk-nonrepetitive coloring  $c \colon V(G) \to \{1, \ldots, k\}$  of G, which is called *the walk-nonrepetitive chromatic number* [Barát and Wood 2008]. Nonrepetitive colorings have attracted some attention in the last years as applications of a method related to the entropy compression method [Rosenfeld 2020]. We also recommend [Wood 2021].

Note that any walk-nonrepetitive coloring c of a graph G is a proper coloring because if  $uv \in E(G)$ , then uv is not a boring walk, and hence  $c(u) \neq c(v)$ . Moreover, all the neighbors of a same vertex must have different colors. Indeed, let  $v \in V(G)$  and uand w be distinct neighbors of v. Since uvwv is not a boring walk, we have  $c(u) \neq c(w)$ . This implies that  $\sigma(G) \ge \Delta(G) + 1$ , where  $\Delta(G)$  denotes that maximum degree of G, and hence  $e(G) \le \frac{1}{2}\Delta(G)|V(G)| \le \frac{\sigma(G)-1}{2}|V(G)|$ , which is the upper bound proved in [Barát and Wood 2008]. This upper bound is tight because if G is a union of disjoint copies of  $K_k$ , then  $\sigma(G) = k$  and  $e(G) = \frac{k-1}{2}|V(G)|$ . We show that such an extremal graph is unique when k divides |V(G)| (Theorem 1), and that a smaller upper bound holds for connected graphs in the case k = 4 (Theorem 2).

### 2. Main results

Let G be a connected graph on n vertices, and note that if  $e(G) = \frac{1}{2}\Delta(G)n$ , then G is *regular*, i.e., all of its vertices have the same degree. The following result implies that, given  $k \ge 4$ , there is a unique graph with walk-nonrepetitive chromatic number k and such a number of edges when k divides n.

**Theorem 1.** Let  $k \ge 4$ . If G is a (k-1)-regular graph for which  $\sigma(G) = k$ , then G is the union disjoint of copies of  $K_k$ .

*Proof.* The proof follows by induction on k. First, suppose that k = 4, and let G be a 3-regular graph for which  $\sigma(G) = 4$ . Let  $C_1, C_2, C_3, C_4$  be the color classes of a minimum walk-nonrepetitive coloring of G. As mentioned above, since  $\sigma(G) = 4$ , and G is 3-regular, each vertex in  $C_i$  has exactly one neighbor in each  $C_j$  with  $j \in \{1, 2, 3, 4\}$  and  $j \neq i$ . Therefore,  $H = G \setminus C_1$  is a 2-regular graph, i.e., a set of disjoint cycles. In fact, since each vertex in H has precisely two neighbors of different colors, one may prove that each component of H is a cycle of length divisible by 3. Note that if one such component C has length at least 6, then each sequence of six consecutive vertices in C forms a repetitive path that is not boring. Therefore, each component of H is a triangle.

Now, we prove that each component of G is a copy of  $K_4$ . Let  $v_1 \in C_1$  and let  $v_2$  and  $v_3$  be distinct neighbors of  $v_1$ . Suppose, for a contradiction, that  $v_2$  and  $v_3$ are in distinct components of H, say  $H_2$  and  $H_3$ , respectively. Suppose, without loss of generality, that  $v_2 \in C_2$  and  $v_3 \in C_3$ , and let  $v'_2$  (resp.  $v'_3$ ) be a neighbor of  $v_3$  (resp.  $v_2$ ) in  $C_2$  (resp.  $C_3$ ). Note that  $v'_2$  and  $v_3$  are in  $H_3$  while  $v'_3$  and  $v_2$  are in  $H_2$ , and hence  $v'_2 \neq v_2$  and  $v'_3 \neq v_3$ . Now, let  $v'_1$  be a neighbor of  $v'_3$  in  $C_1$ , and note that  $v'_1v'_3v_2v_1v_3v'_2$ is a repetitive path that is not boring, a contradiction. Therefore all of the neighbors of  $v_1$ are in the same component of H, and hence H is a union of disjoint copies of  $K_4$ . Now, suppose k > 4. Analogously to the case above,  $H = G \setminus C_1$  is a (k - 2)-regular graph for which  $\sigma(H) = k - 1$ . By the induction hypothesis, H is a union of disjoint copies of  $K_{k-1}$ , and analogously to the case above, all the neighbors of a vertex in  $C_1$  must be in the same component, as desired.

Theorem 1 implies that any connected graph G for which  $|V(G)| \ge 5$  and  $\sigma(G) = 4$  has less than  $\frac{3}{2}|V(G)|$  edges. The next result slightly improves this upper bound.

**Theorem 2.** If G is a connected graph on  $n \ge 5$  vertices for which  $\sigma(G) = 4$ , then  $e(G) \le 4n/3$ .

*Proof.* The proof follows by contradiction. Let G be a connected graph with  $n \ge 5$  vertices and  $\sigma(G) = 4$ , and suppose that e(G) > 4n/3. Fix a minimum walk-nonrepetitive coloring c of G. The first step of the proof is to show that G does not have some special subgraphs. For example, G has no copies of  $K_4$ . Indeed, since  $\Delta(G) \le 3$ , if G has a copy K of  $K_4$ , then K is a component of G, and hence G = K, a contradiction to n > 4.

In what follows, let  $V_3$  be the set of vertices of G with degree exactly 3, let  $V_{\leq 2} = V(G) \setminus V_3$ , and set  $G_3 = G[V_3]$ . Now, let Y be the tree obtained from the star with three edges by subdividing one of its edges (see Figure 1). We claim that  $G_3$  has no copies of Y as a subgraph. Indeed, suppose that  $G_3$  has a copy H of Y on the vertices  $u_1, u_2, u_3, u_4, w$ , where  $N_H(u_1) = \{u_2, u_3, u_4\}$  and  $N_H(u_4) = \{u_1, w\}$ . Then  $u_2, u_3$  and w are leaves of H. We may suppose, without loss of generality, that  $c(u_i) = i$ , for every  $i \in \{1, 2, 3, 4\}$ , and hence c(w) = 2 or c(w) = 3. Say c(w) = 2. The case c(w) = 3 follows analogously. As  $w, u_2 \in V_3$ , we have  $d(w) = d(u_2) = 3$ , and hence w has a neighbor  $v_1$  for which  $c(v_1) = 1$ , and  $u_2$  has a neighbor  $v_4$  for which  $c(v_4) = 4$ . Note that  $v_1 \neq u_1$ , otherwise  $d(u_1) \geq 4$ . Then  $v_1wu_4u_1u_2v_4$  is a repetitive walk that is not boring, a contradiction.



Figure 1. The Y graph.

Figure 2. The R graph.

Now, let R be graph obtained from a  $K_4$  by removing an edge  $v_2v_3$ , and adding two new vertices, each adjacent to one of  $v_2$  and  $v_3$  (see Figure 2). We claim that Ghas no copies of R as a subgraph. Indeed, suppose that G has a copy H of R on the vertices  $v_1, v_2, v_3, v_4, v_5, v_6$ , where  $N_H(v_5) = \{v_2\}$ ,  $N_H(v_6) = \{v_3\}$ . We may suppose that  $c(v_i) = i$ , for every  $i \in \{1, 2, 3, 4\}$ , and hence  $c(v_5) = c(v_3) = 3$  and  $c(v_6) = c(v_2) =$ 2. If  $d_G(v_5) = d_G(v_6) = 1$ , then G = H, and hence e(G) = 7 < 8 = 24/3 = 4n/3, a contradiction. Then either  $v_5$  or  $v_6$ , say  $v_5$ , has a neighbor, say w, outside H. Since cis a walk-nonrepetitive coloring,  $c(w) \notin \{c(v_2), c(v_5)\} = \{2, 3\}$ , and hence  $c(w) = i \in \{1, 4\}$ . Then  $wv_5v_2v_iv_3v_6$  is a repetitive walk that is not boring, a contradiction. Given two disjoint sets of vertices U and W, we denote by e(U, W) the number of edges joining vertices of U to vertices of W. Now, let  $\mathcal{H}$  be the set of components of  $G_3$ , and given  $H \in \mathcal{H}$ , let  $e_H = e(V(H), V_{\leq 2})$ .

**Claim 1.** For every component H of  $G_3$  we have  $|V(H)| \le e_H$ .

*Proof.* If  $\Delta(H) \leq 2$ , then each vertex of H has a neighbor in  $V_{\leq 2}$  and the claim follows. Thus, we may assume that H has a vertex u for which  $d_H(u) = 3$ . Let  $v_1, v_2, v_3$  be the neighbors of u. If  $v_i$  has a neighbor w in  $V_3 \setminus \{u, v_1, v_2, v_3\}$ , then  $\{u, v_1, v_2, v_3, w\}$  induces a copy of Y in  $G_3$ , a contradiction. Thus H has only four vertices and hence  $e_H = 6 - 2e(G[\{v_1, v_2, v_3\}])$ . If  $e(G[\{v_1, v_2, v_3\}]) \leq 1$ , then the claim follows, and if  $e(G[\{v_1, v_2, v_3\}]) = 3$ , then H is a copy of  $K_4$ , which is a contradiction. Then  $e(G[\{v_1, v_2, v_3\}]) = 2$ , and H is the graph obtained from  $K_4$  by removing one edge. Suppose that  $v_1v_2 \notin E(H)$  and let  $w_1$  and  $w_2$  be the remaining neighbors of  $v_1$  and  $v_2$ , respectively. Since G does not have a copy of R, we have  $w_1 = w_2$ . Note that  $c(w_1) \notin \{c(v_1), c(v_2)\}$  because c is a proper coloring. Then  $c(w_1) \in \{c(u), c(v_3)\}$ , and hence  $v_1$  has two neighbors with the same color, a contradiction.

By Claim 1, we have  $|V_3| = \sum_{H \in \mathcal{H}} |V(H)| \le \sum_{H \in \mathcal{H}} e_H = e(V_3, V_{\le 2})$ . Moreover,  $e(V_3, V_{\le 2}) \le 2|V_{\le 2}|$ , because  $d(v) \le 2$  for every  $v \in V_{\le 2}$ . Therefore,  $|V_3| \le 2|V_{\le 2}|$ . Since  $|V_3| + |V_{\le 2}| = n$ , we have  $|V_3| \le 2n/3$ . Therefore,

$$2e(G) = \sum_{v \in V_3} d(v) + \sum_{v \in V_{\leq 2}} d(v) \leq 3|V_3| + 2|V_{\leq 2}| = 2n + |V_3| \leq 2n + \frac{2n}{3} = \frac{8n}{3}$$

as desired.

### 3. Future work

In this paper we have shown the uniqueness of the extremal graph with walk-nonrepetitive chromatic number k and order divisible by k for any fixed  $k \ge 4$ , which is a disconnected graph; and we also presented a smaller upper bound on the maximum number of edges in a connected graph G with walk-nonrepetitive chromatic number 4. So far we have not found a graph with a matching number of edges, so this upper bound may still be improved. We were also able to show that, for general  $k \ge 4$ , G does not contain generalization of the Y graph, but a stronger structure seems to be needed to generalize this bound. These results may be later combined to characterize the extremal graphs with any order.

#### References

Barát, J. and Wood, D. R. (2008). Notes on nonrepetitive graph colouring. *The Electronic Journal of Combinatorics*, 15, R99.

- Rosenfeld, M. (2020). Another approach to non-repetitive colorings of graphs of bounded degree. *The Electronic Journal of Combinatorics*, 27(3), P3.43.
- Wood, D. R. (2021). Nonrepetitive graph colouring. *The Electronic Journal of Combina*torics, DS24.